

Momentum

Defining MOMENTUM

Momentum \equiv Mass x Velocity

Momentum is given by the symbol: \vec{p} . So, $\vec{p} = m\vec{v}$

You can think of momentum as the “amount of motion” something has.

For example, which has a momentum of greater magnitude?

A) A 1000 kg car travelling at 60 mi/hr, or

B) A 20000 kg locomotive travelling at 3 mi/hr.

Answer: Both have the same magnitude of momentum.

Defining IMPULSE

Impulse is a CHANGE in momentum:

$$\Delta\vec{p} = \vec{p}_{final} - \vec{p}_{initial}$$

Recall that $\vec{F}_{net} = m\vec{a}$ and that $\vec{a} = \frac{\Delta\vec{v}}{\Delta t}$. Then $\vec{F}_{net} = m\frac{\Delta\vec{v}}{\Delta t}$.

Thus, $\vec{F}_{net}\Delta t = m\Delta\vec{v} = m\vec{v}_{final} - m\vec{v}_{initial} = \vec{p}_{final} - \vec{p}_{initial} = \Delta\vec{p}$

Standard units of momentum and impulse:

Mass Units x Velocity Units = (kilograms) x (meters/second) = **kg·m/s** OR

Force Units x Time Units = (Newtons)x(seconds) = **N·s**

Momentum & Impulse are VECTORS

Both momentum and impulse have directional aspects. They are vectors and are fully described by having BOTH magnitude AND direction.

Compare the momentum of a 20 kg cheetah sprinting at 25 m/s to that of a 25 kg cheetah sprinting at 20 m/s.

- A) They both have the same momentum.
- B) Not enough information given.

Answer: (B) Not enough information...


(We don't know the directions the cheetahs are sprinting.)

Example involving Impulse

A 0.2-kg ball of clay ball is dropped. The clay ball hits the floor at a speed of 8.0 m/s and sticks to the floor. What impulse did the floor impart to the ball to change its momentum?

$$\begin{aligned}\text{Impulse} &= \Delta\vec{p} = \vec{p}_{final} - \vec{p}_{initial} \\ &= m\vec{v}_{final} - m\vec{v}_{initial} \\ &= (0.2 \text{ kg})(0 \text{ m/s}) - (0.2 \text{ kg})(-8.0 \text{ m/s}) \\ &= +1.6 \text{ kg}\cdot\text{m/s} \\ &\text{(The “+” sign implying “upward.”)}\end{aligned}$$

$$\text{○ } m = 0.2 \text{ kg}$$




Follow-up to the previous example:

A 0.2-kg rubber ball is dropped. The rubber ball hits the floor at a speed of 8.0 m/s and rebounds back up at the same speed. What impulse did the floor impart to the ball to change its momentum?

$$\begin{aligned}\text{Impulse} &= \Delta \vec{p} = \vec{p}_{final} - \vec{p}_{initial} \\ &= m\vec{v}_{final} - m\vec{v}_{initial} \\ &= (0.2 \text{ kg})(+8.0 \text{ m/s}) - (0.2 \text{ kg})(-8.0 \text{ m/s}) \\ &= +3.2 \text{ kg}\cdot\text{m/s}\end{aligned}$$

(Note that it takes a greater impulse to make something bounce backward.)



Conservation of Momentum

What makes momentum a useful concept?


If the net force on a system is zero, then $\vec{F}_{net}\Delta t = 0$ and there is no impulse imparted to the system. Therefore, the momentum of the system remains constant. This is conservation of momentum.

Consider cue ball (mass m) hitting a stationary 8-ball (same mass m):



By conservation of \vec{p} , $m\vec{v} + 0 = 0 + m\vec{v}$, so we know that the velocity of the 8-ball after the collision is the same as the incoming cue ball which comes to rest.

Consider a two-cart system where a moving cart bumps into a stationary cart. After the collision, the two carts stick together. Calculate the speed v_{final} of the combination after the collision.



Use Conservation of Momentum: $\vec{p}_{initial} = \vec{p}_{final}$

$$m_1 \vec{v}_{1,initial} + m_2 \vec{v}_{2,initial} = (m_1 + m_2) \vec{v}_{final}$$


$$\vec{v}_{final} = \frac{m_1 \vec{v}_{1,initial}}{(m_1 + m_2)}$$

2-Dimensional Collisions

The figure shows an overhead view of two cars of the same mass approaching an intersection where they collide and stick together.

The speed of the northbound car is twice that of the eastbound car.

What path does the wreckage take?



Answer: Path B